




## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>5</sup> : <b>A61L 9/16</b>		A1	(11) International Publication Number: <b>WO 94/23767</b>
			(43) International Publication Date: <b>27 October 1994 (27.10.94)</b>
(21) International Application Number: <b>PCT/US94/04560</b>		(81) Designated States: AT, AU, BB, BG, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KG, KP, KR, KZ, LK, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: <b>22 April 1994 (22.04.94)</b>			
(30) Priority Data: 051,019      22 April 1993 (22.04.93)      US 205,511      2 March 1994 (02.03.94)      US			
(71) Applicant (for all designated States except US): <b>EMISPHERE TECHNOLOGIES, INC. [US/US]; 15 Skyline Drive, Hawthorne, NY 10532 (US).</b>		<b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(72) Inventors; and			
(75) Inventors/Applicants (for US only): <b>MILSTEIN, Sam, J. [US/US]; 1 Knollwood Drive, Larchmont, NY 10538 (US). BARANTSEVITCH, Evgueni, N. [RU/US]; Apartment 2B, 127 Cooper Drive, New Rochelle, NY 10801 (US). SARUBBI, Donald, J. [US/US]; 18 Lawton Lane, Bronxville, NY 10708 (US). LEONE-BAY, Andrea [US/US]; 20 Woodland Way, Ridgefield, CT 06877 (US). PATON, Duncan, R. [GB/US]; 323 Nash Road, Purdys, NY 10378 (US).</b>			
(74) Agents: <b>ROBINSON, Joseph, R. et al.; Darby &amp; Darby, 805 Third Avenue, New York, NY 10022 (US).</b>			

(54) Title: ORAL DRUG DELIVERY COMPOSITIONS AND METHODS



## (57) Abstract

The present invention relates to an oral drug delivery system, and in particular to modified amino acids and modified amino acid derivatives for use as a delivery system of sensitive agents such as bioactive peptides. The modified amino acids and derivatives can form non-covalent mixtures with active biological agents and in an alternate embodiment can releasably carry active agents. Modified amino acids can also form drug containing microspheres. These mixtures are suitable for oral administration of biologically active agents to animals. Methods for the preparation of such amino acids are also disclosed.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

5

10

**ORAL DRUG DELIVERY COMPOSITIONS AND METHODS**

This application is a continuation-in-part of U.S. Patent Application Serial No. 08/051,019, filed on April 22, 1993 and U.S. Patent Application Serial No. 08/205,511 filed on March 2, 1994.

**FIELD OF THE INVENTION**

The present invention relates to compositions suitable for oral drug delivery, and in particular to compositions in which modified amino acids and modified amino acid derivatives are used as carriers for sensitive agents such as bioactive peptides and the like. The modified amino acids or derivatives can form non-covalent mixtures with biologically-active agents which are suitable for oral administration to animals. Methods for the preparation and for the administration of such compositions are also disclosed.

**Background of the Invention**

Conventional means for delivering biologically-active agents, including, but not limited to, pharmaceutical and therapeutic agents, to animals are often severely limited by chemical barriers and physical barriers imposed by the body. Oral delivery of many biologically-active agents would be the route of choice if not for chemical and physico-chemical barriers such as the extreme and varying pH in the gastro-intestinal (GI) tract, exposure to powerful digestive enzymes, and the impermeability of gastro-intestinal membranes to the active agent. Among the numerous agents which are not typically suitable for oral administration are biologically-active peptides such as

calcitonin and insulin. Examples of other compounds which are affected by these physico-chemical barriers are polysaccharides and particularly mucopolysaccharides, including, but not limited to, heparin; heparinoids; 5 antibiotics; and other organic substances. These agents are rapidly destroyed in the gastro-intestinal tract by acid hydrolysis, enzymes, or the like.

Prior methods for orally administering vulnerable pharmacological agents have relied on the co-administration 10 of adjuvants (e.g., resorcinols and non-ionic surfactants such as polyoxyethylene oleyl ether and n-hexadecyl polyethylene ether) to increase artificially the permeability of the intestinal walls; and on the co-administration of enzymatic inhibitors (e.g., pancreatic 15 trypsin inhibitor, diisopropylfluorophosphate (DFF) and trasylol) to inhibit enzymatic degradation. Liposomes have also been described as drug delivery systems for insulin and heparin. See, for instance, U.S. Patent No. 4,239,754; Patel et al. (1976) FEBS Letters Vol. 62, page 60; and Hashimoto 20 et al. (1979) Endocrinol. Japan, Vol. 26, page 337.

However, broad spectrum use of the aforementioned drug delivery systems is precluded for reasons including: (1) the need to use toxic amounts of adjuvants or inhibitors; (2) the lack of suitable low MW cargoes; (3) the poor stability 25 and inadequate shelf life of the systems; (4) the difficulties in manufacturing the systems; (5) the failure of the systems to protect the active ingredient; and (6) the failure of the systems to promote absorption of the active agent.

30 More recently, microspheres of artificial polymers or proteinoids of mixed amino acids have been described for delivery of pharmaceuticals. For example, U.S. Patent No. 4,925,673 describes drug containing microsphere constructs as well as methods for their preparation and use. These 35 proteinoid microspheres are useful for delivery of a number of active agents.

There is still a need in the art for simple and inexpensive delivery systems which are easily prepared and


which can deliver a broad range of biologically-active agents.

### Summary of the Invention

5 Compositions for orally delivering biologically-active agents incorporating modified amino acids, amino acid derivatives, peptides and peptide derivatives as carriers are provided. These compositions comprise

- (A) at least one biologically-active agent; and
- 10 (B) at least one carrier comprising
  - (a) (i) at least one acylated aldehyde of an amino acid,
  - (ii) at least one acylated ketone of an amino acid,
  - 15 (iii) at least one acylated aldehyde of a peptide,
  - (iv) at least one acylated ketone of a peptide, or
  - (v) any combination of (a)(i), (a)(ii),
  - 20 (a)(iii) and (a)(iv);

- (b) (i) carboxymethyl-phenylalanine-leucine,
- (ii) 2-carboxy-3-phenylpropionyl-leucine,
- (iii) 2-benzylsuccinic acid,
- 25 (iv) an actinonin, or
- (v) a compound having the formula:



wherein: Ar is a substituted or unsubstituted phenyl or naphthyl;

30 Y is  $\text{--}\overset{\text{O}}{\parallel}\text{C--}$  or  $\text{--SO}_2\text{--}$ ;

$\text{R}^1$  is  $\text{--N}(\text{R}^4)\text{--R}^3\text{--}\overset{\text{O}}{\parallel}\text{C--}$ ; wherein:

35  $\text{R}^3$  is  $\text{C}_1$  to  $\text{C}_{24}$  alkyl,  $\text{C}_1$  to  $\text{C}_{24}$  alkenyl, phenyl, naphthyl, ( $\text{C}_1$  to  $\text{C}_{10}$  alkyl)phenyl, ( $\text{C}_1$  to  $\text{C}_{10}$  alkenyl)phenyl, ( $\text{C}_1$  to  $\text{C}_{10}$  alkyl)naphthyl, ( $\text{C}_1$  to  $\text{C}_{10}$  alkenyl)naphthyl, phenyl( $\text{C}_1$  to  $\text{C}_{10}$  alkyl), phenyl( $\text{C}_1$  to  $\text{C}_{10}$  alkenyl), naphthyl( $\text{C}_1$  to  $\text{C}_{10}$  alkyl) and naphthyl( $\text{C}_1$  to  $\text{C}_{10}$  alkenyl);

R<sup>3</sup> is optionally substituted with C<sub>1</sub> to C<sub>4</sub> alkyl, C<sub>1</sub> to C<sub>4</sub> alkenyl, C<sub>1</sub> to C<sub>4</sub> alkoxy, -OH, -SH, -CO<sub>2</sub>R<sup>5</sup>, cycloalkyl, cycloalkenyl, heterocyclic, aryl, alkaryl, heteroaryl or heteroalkaryl or any combination thereof;

5 R<sup>5</sup> is hydrogen, C<sub>1</sub> to C<sub>4</sub> alkyl or C<sub>1</sub> to C<sub>4</sub> alkenyl;

R<sup>3</sup> is optionally interrupted by oxygen, nitrogen, sulfur or any combination thereof; and

10 R<sup>4</sup> is hydrogen, C<sub>1</sub> to C<sub>4</sub> alkyl or C<sub>1</sub> to C<sub>4</sub> alkenyl;

and n is from 1 to about 5;

(vi) or any combination of (b) (i), (b) (ii),

(b) (iii), (b) (iv) and (b) (v); or

(c) a combination of (a) and (b).

15 Also contemplated is a method for preparing these compositions which comprises mixing at least one biologically active agent, with at least one carrier as described above, and optionally, a dosage vehicle.

In an alternative embodiment, these non-toxic  
20 carriers are orally administered to animals as part of a delivery system by blending or mixing the carriers with a biologically active agent prior to administration. The carriers may also form microspheres in the presence of the active agent. The microspheres containing the active agent  
25 are then orally administered. Also contemplated by the present invention are dosage unit forms that include these compositions.

#### Description of the Drawings

30 Figure 1 is a graphic illustration of the results of oral gavage testing in rats using salmon calcitonin with acetyl phenylalanine aldehyde, carbomethoxyPhe-Leu-OH, and acetyl-Phe-Leu-Leu-Arg aldehyde carriers.

35 Figure 2 is a graphic illustration of the results of oral gavage testing in rats using salmon calcitonin with acetylleucine aldehyde and acetylphenylalanine aldehyde carriers.

Figure 3 is a graphic illustration of the results of oral gavage testing in rats using salmon calcitonin with acetylphenylalanine aldehyde and carbomethoxyPhe-Leu-OH carriers.

5           Figure 4 is a graphic illustration of the results of oral gavage testing in rats using salmon calcitonin with acetylphenylalanine aldehyde, acetylLeu-Leu-Arg aldehyde and carbomethoxyPhe-Leu-OH carriers.

10           Figure 5 is a graphic illustration of the results of intraduodenal injection testing in rats using salmon calcitonin with acetylphenylalanine aldehyde and 4-(phenylsulfonamido)-4-phenylbutyric acid carriers.

15           Figure 6 is a graphic illustration of the results of oral gavage testing in rats using salmon calcitonin with acetylphenylalanine aldehyde, N-acetyllysine, and acetyl-Leu aldehyde carriers.

20           Figure 7 is a graphic illustration of the results of intraduodenal injection testing in rats using salmon calcitonin with acetylphenylalanine aldehyde carrier in aqueous ethanol, dimethyl sulfoxide (DMSO), and olive oil dosing vehicles, and in a DMSO dosing vehicle alone.

Figure 8 is a graphic illustration of the results of oral gavage testing in rats using salmon calcitonin with cyclohexanoyl-phenylalanine aldehyde carrier.

25           Figure 9 is a graphic illustration of rat serum calcium levels after oral administration of two dosage levels of a modified amino acid microsphere preparation containing salmon calcitonin and a soluble modified amino acid preparation containing salmon calcitonin after pre-  
30 dosing with a sodium bicarbonate solution.

Figure 10 is a graphic illustration of the results of oral gavage testing in rats using salmon calcitonin with acetyl-Phe aldehyde, actinonin, and carbomethoxy-Phe-Leu-OH carriers.

35           Figure 11 is a graphic illustration of the results of oral gavage testing in rats using salmon calcitonin with 4-(phenylsulfonamido)-4-phenylbutyric acid carrier.

Figure 12 is a graphic illustration of the results of oral gavage testing in rats using salmon calcitonin with 3-(phenylsulfonamido)benzoic acid and 4-(phenylsulfonamido)-hippuric acid carriers.

5           Figure 13 is a graphic illustration of the results of oral gavage testing in rats using salmon calcitonin with 4-(phenylsulfonamido)-4-phenylbutyric acid and 4-(phenylsulfonamido)benzoic acid carriers.

10           Figure 14 is a graphic illustration of the results of oral gavage testing in rats using salmon calcitonin with 4-(phenylsulfonamido)-4-phenylbutyric acid and 4-(phenylsulfonamido)phenylacetic acid carriers.

15           Figure 15 is a graphic illustration of the results of oral gavage testing in rats using interferon  $\alpha 2b$  (rhIFN) with 4-(phenylsulfonamido)-4-phenylbutyric acid carrier.

            Figure 16 is a graphic illustration of the results of oral gavage testing in rats using interferon  $\alpha 2b$  with 4-(phenylsulfonamidomethyl)benzoic acid carrier.

20           Figure 17 is a graphic illustration of the results of oral gavage testing in rats using interferon  $\alpha 2b$  with 4-(phenylsulfonamido)phenylacetic acid as carrier.

            Figure 18 is a graphic illustration of the results of oral gavage testing in rats using interferon  $\alpha 2b$  with 4-(phenylsulfonamido)hippuric acid carrier.

25           Figures 19 and 20 are graphic illustrations of the results of oral gavage testing in hypophysectomized rats using growth hormone alone and at two dosage levels with 4-(phenylsulfonamido)-4-phenylbutyric acid carrier.

30           Figure 21 is a graphic illustration of the results of oral gavage testing in normal rats using growth hormone with 4-(phenylsulfonamido)-4-phenylbutyric acid carrier.

            Figure 22 is a graphic illustration of the results of oral gavage testing in rats using disodium cromoglycate with 4-(phenylsulfonamido)-4-phenylbutyric acid as carrier.

35

#### Detailed Description of the Invention

Amino acids and amino acid derivatives, in modified form, may be used to deliver orally sensitive



biologically-active agents, including, but not limited to, hormones such as calcitonin, insulin, and polysaccharides such as heparin, which would not be considered orally administrable for various reasons. Insulin, for example is sensitive to the denaturing conditions of the gastro-intestinal (GI) tract. Also, heparin, by virtue of its charge and hydrophilic nature, is not readily absorbed from the gastro-intestinal tract. In contrast to the modified amino acids and modified amino acid derivatives of the present invention, unmodified free amino acids do not provide protection against degradation in the GI tract for labile bioactive agents.

The compositions of the subject invention are useful for administering biologically-active agents to any animals such as birds; mammals, such as primates and particularly humans; and insects.

Other advantages provided by the present invention include the use of readily available and inexpensive starting materials in cost-effective methods for preparing and isolating modified amino acid derivatives. These methods are simple to perform and are amenable to industrial scale-up for commercial production.

Biologically-active agents suitable for use with carriers disclosed herein include, but are not limited to, peptides, and particularly small peptide hormones, which by themselves do not pass or only pass slowly through the gastro-intestinal mucosa and/or are susceptible to chemical cleavage by acids and enzymes in the gastro-intestinal tract; polysaccharides and particularly mixtures of mucopolysaccharides; carbohydrates; lipids; or any combination thereof. Examples include, but are not limited to, human growth hormone; bovine growth hormone; growth hormone releasing hormone; interferons; interleukin-I; insulin; heparin, and particularly low molecular weight heparin; calcitonin; erythropoietin; atrial natriuretic factor; antigens; monoclonal antibodies; somatostatin; adrenocorticotropin; gonadotropin releasing hormone; oxytocin; vasopressin; cromolyn sodium (sodium or disodium

cromoglycate); vancomycin; desferrioxamine (DFO); or any combination thereof.

Additionally the carriers of the present invention can be used to deliver other active agents such as  
5 pesticides and the like.

The term amino acid as used herein includes any carboxylic acid having at least one free amine group including naturally occurring and synthetic amino acids. The preferred amino acids are  $\alpha$ -amino acids, and preferably  
10 are naturally occurring  $\alpha$ -amino acids although non- $\alpha$ -amino acids are useful as well.

Poly amino acids as used herein refers to peptides or two or more amino acids linked by a bond formed by other groups which can be linked, e.g., an ester, anhydride or an  
15 anhydride linkage.

The term peptide is meant to include two or more amino acids joined by a peptide bond. Peptides can vary in length from dipeptides with 2 to poly peptides with several hundred amino acids. See Chambers Biological Dictionary,  
20 editor Peter M. B. Walker, Cambridge, England: Chambers Cambridge, 1989, page 215. The peptides most useful in the practice of the present invention include di-peptides, tri-peptides, tetra-peptides, and penta-peptides. The preferred peptides are di-peptides, tri-peptides. Peptides  
25 can be homo- or hetero- peptides and can include natural amino acids, synthetic amino acids, or any combination thereof.

The term amino acid derivatives and peptide derivatives as used herein are meant to include amino acid  
30 aldehydes or ketones and/or peptide aldehydes or ketones where the -COOH group has been converted to a ketone or aldehyde.

The terms modified amino acids, peptides, and derivatives thereof are meant to include amino acids, amino  
35 acid derivatives, peptides and peptide derivatives which have been modified as described below by acylating or sulfonating at least one free amine group, with an acylating

or sulfonating agent which reacts with at least one of the free amine groups present.

The preferred naturally occurring amino acids for use in the present invention as amino acids or components of a peptide are alanine, arginine, asparagine, aspartic acid, citrulline, cysteine, cystine, glutamine, glycine, histidine, isoleucine, leucine, lysine, methionine, ornithine, phenylalanine, proline, serine, threonine, tryptophan, tyrosine, valine, hydroxy proline,  $\gamma$ -carboxyglutamate, or O-phosphoserine. The most preferred amino acids are arginine, leucine, lysine, phenylalanine, tyrosine and valine.

The preferred non-naturally occurring amino acids for use in the present invention as amino acids or components of a peptide are  $\beta$ -alanine, phenylglycine,  $\alpha$ -aminobutyric acid,  $\gamma$ -amino butyric acid, 4-(4-aminophenyl)butyric acid,  $\alpha$ -amino isobutyric acid,  $\epsilon$ -aminocaproic acid, 7-aminoheptanoic acid,  $\beta$ -aspartic acid, aminobenzoic acid, (aminomethyl)benzoic acid, aminophenylacetic acid, aminohippuric acid,  $\gamma$ -glutamic acid, cysteine(ACM),  $\epsilon$ -lysine,  $\epsilon$ -lysine (A-Fmoc), methionine sulfone, norleucine, norvaline, ornithine, d-ornithine, p-nitrophenylalanine, hydroxy proline, and thioproline.

The amino acids useful in the practice of the subject invention have the formula:



$\text{R}^2$  has the formula  $-\text{R}^3 - \overset{\text{O}}{\parallel}{\text{C}} -$  wherein  $\text{R}^3$  is  $\text{C}_1$  to  $\text{C}_{24}$  alkyl,  $\text{C}_1$  to  $\text{C}_{24}$  alkenyl, phenyl, naphthyl, ( $\text{C}_1$  to  $\text{C}_{10}$  alkyl)-phenyl, ( $\text{C}_1$  to  $\text{C}_{10}$  alkenyl)phenyl, ( $\text{C}_1$  to  $\text{C}_{10}$  alkyl)naphthyl, ( $\text{C}_1$  to  $\text{C}_{10}$  alkenyl)naphthyl, phenyl( $\text{C}_1$  to  $\text{C}_{10}$  alkyl), phenyl( $\text{C}_1$  to  $\text{C}_{10}$  alkenyl), naphthyl( $\text{C}_1$  to  $\text{C}_{10}$  alkyl) and naphthyl( $\text{C}_1$  to  $\text{C}_{10}$  alkenyl);

optionally  $\text{R}^3$  is substituted with  $\text{C}_1$  to  $\text{C}_4$  alkyl,  $\text{C}_1$  to  $\text{C}_4$  alkenyl,  $\text{C}_1$  to  $\text{C}_4$  alkoxy, -OH, -SH and  $-\text{CO}_2\text{R}^5$  or any combination thereof;

$\text{R}^5$  is hydrogen,  $\text{C}_1$  to  $\text{C}_4$  alkyl or  $\text{C}_1$  to  $\text{C}_4$  alkenyl;

$R^3$  is optionally interrupted by oxygen, nitrogen, sulfur or any combination thereof; and

$R^4$  is hydrogen,  $C_1$  to  $C_4$  alkyl or  $C_1$  to  $C_4$  alkenyl.

The phenyl or naphthyl groups can be optionally substituted. Suitable but non-limiting examples of substitutents are  $C_1$  to  $C_6$  alkyl,  $C_1$  to  $C_6$  alkenyl, alkoxy having from 1 to 6 carbon atoms, hydroxy, thio, or  $CO_2R^6$  wherein  $R^6$  is hydrogen,  $C_1$  to  $C_6$  alkyl,  $C_1$  to  $C_6$  alkenyl.

The amino acid derivatives or peptide derivatives of the present invention can be readily prepared by reduction of amino acid esters or peptide esters with an appropriate reducing agent. For example, amino acid aldehydes or peptide aldehydes can be prepared as described in an article by R. Chen et al., *Biochemistry*, 1979, 18, 921-926. Amino acid or peptide ketones can be prepared by the procedure described in Organic Syntheses, Col. Vol. IV, Wiley, (1963), pages 5-6. Amino acids, peptides, amino acid esters, peptide esters, and other necessary reagents to prepare these derivatives are readily available from a number of commercial sources such as Aldrich Chemical Co. (Milwaukee, WI, USA); Sigma Chemical Co. (St. Louis, MO, USA); and Fluka Chemical Corp. (Ronkonkoma, NY, USA).

The amino acids and peptides are modified by acylating or sulfonating at least one free amine group, with an acylating or sulfonating agent which reacts with at least one of the free amine groups present. Suitable, but non-limiting, examples of agents useful for modifying amino acids or peptides useful in practicing the present invention include

acylating and sulfonating agents having the formula  $R^7-\overset{\text{O}}{\underset{\text{||}}{\text{C}}}-X$  or  $R^7-\text{SO}_2-X$  wherein  $R^7$  is alkyl or alkenyl, preferably having from 1 to 20 carbon atoms, or aromatic preferably having from 6 to 20 carbon atoms.

The  $R^7$  group can be substituted or unsubstituted. The preferred substitutents include  $C_1$  to  $C_4$  alkyl,  $C_1$  to  $C_4$  alkenyl,  $C_1$  to  $C_4$  alkoxy,  $CO_2R^8$  wherein  $R^8$  is hydrogen,  $C_1$  to  $C_4$  alkyl or  $C_1$  to  $C_4$  alkenyl.

Preferably,  $R^7$  is methyl, ethyl, phenyl, benzyl or naphthyl. More preferably,  $R^7$  is phenyl, or acetyl. X is a leaving group. In a reaction in which the substrate molecule becomes cleaved, part of it (the part not

5 containing the carbon) is usually called the *leaving group*. See Advanced Organic Chemistry, 2d edition, Jerry March, New York: McGraw-Hill Book (1977), page 187, Typical leaving groups include, but are not limited to, halogens such as chlorine, bromine and iodine.

10 Examples of the acylating and sulfonating agents for amino acids and peptides include, but are not limited to, acyl halides such as acetyl chloride, propyl chloride, benzoyl chloride, hippuryl chloride and the like; sulfonyl halides such as benzene sulfonyl chloride, and anhydrides,  
15 such as acetic anhydride, propyl anhydride, benzoic anhydride, hippuric anhydride and the like. The preferred acylating and sulfonating agents are benzoyl chloride, benzene sulfonyl chloride, and hippuryl chloride.

The modified acid compounds have the formula:

20 
$$\text{Ar}-\text{Y}-(\text{R}^1)_n-\text{OH}$$

wherein Ar is a substituted or unsubstituted phenyl or naphthyl;

Y is  $\text{—}\overset{\text{O}}{\parallel}\text{C—}$  or  $\text{—SO}_2\text{—}$ ,  $\text{R}^1$  has the formula  $\text{—N}(\text{R}^4)\text{—R}^3\text{—}\overset{\text{O}}{\parallel}\text{C—}$ , wherein:

25  $\text{R}^3$  is  $\text{C}_1$  to  $\text{C}_{24}$  alkyl,  $\text{C}_1$  to  $\text{C}_{24}$  alkenyl, phenyl, naphthyl, ( $\text{C}_1$  to  $\text{C}_{10}$  alkyl) phenyl, ( $\text{C}_1$  to  $\text{C}_{10}$  alkenyl) phenyl, ( $\text{C}_1$  to  $\text{C}_{10}$  alkyl) naphthyl, ( $\text{C}_1$  to  $\text{C}_{10}$  alkenyl) naphthyl, phenyl ( $\text{C}_1$  to  $\text{C}_{10}$  alkyl), phenyl ( $\text{C}_1$  to  $\text{C}_{10}$  alkenyl), naphthyl ( $\text{C}_1$  to  $\text{C}_{10}$  alkyl) and naphthyl ( $\text{C}_1$  to  $\text{C}_{10}$  alkenyl);

30  $\text{R}^3$  is optionally substituted with  $\text{C}_1$  to  $\text{C}_4$  alkyl,  $\text{C}_1$  to  $\text{C}_4$  alkenyl,  $\text{C}_1$  to  $\text{C}_4$  alkoxy,  $\text{—OH}$ ,  $\text{—SH}$  and  $\text{—CO}_2\text{R}^5$  or any combination thereof;

$\text{R}^5$  is hydrogen,  $\text{C}_1$  to  $\text{C}_4$  alkyl or  $\text{C}_1$  to  $\text{C}_4$  alkenyl;

35  $\text{R}^3$  is optionally interrupted by oxygen, nitrogen, sulfur or any combination thereof; and

$\text{R}^4$  is hydrogen,  $\text{C}_1$  to  $\text{C}_4$  alkyl or  $\text{C}_1$  to  $\text{C}_4$  alkenyl.

The amino acid derivatives and peptide derivatives are modified by acylating at least one free amine group,

with an acylating agent which reacts with at least one of the free amine groups present. Suitable, but non-limiting, examples of acylating agents useful for modifying amino acid derivatives and peptide derivatives useful in practicing the present invention include

O


acid chloride acylating agents having the formula  $R^9-C-X$  or wherein:

$R^9$  is alkyl or alkenyl, preferably having from 1 to 20 carbon atoms, cycloalkyl or cycloalkenyl, preferably having from 1 to 20 carbon atoms, or aromatic preferably having from 6 to 20 carbon atoms. The  $R^9$  group can be substituted or unsubstituted. The preferred substituents include  $C_1$  to  $C_4$  alkyl,  $C_1$  to  $C_4$  alkenyl,  $C_1$  to  $C_4$  alkoxy,  $CO_2R^{10}$  wherein  $R^{10}$  is hydrogen,  $C_1$  to  $C_4$  alkyl or  $C_1$  to  $C_4$  alkenyl. Preferably,  $R^9$  is methyl, ethyl, cyclohexyl, cyclopentyl, cycloheptyl, phenyl, benzyl or naphthyl. More preferably,  $R^9$  is phenyl, cyclohexyl cyclopentyl, cycloheptyl, or acetyl. X is a leaving group. Typical leaving groups include, but are not limited to, halogens such as chlorine, bromine and iodine.

Examples of the acylating agents for amino acid derivatives and peptide derivatives include, but are not limited to, acyl halides such as acetyl chloride, propyl chloride, cyclohexanoyl chloride, cyclopentanoyl chloride, and cycloheptanoyl chloride, benzoyl chloride, hippuryl chloride and the like; and anhydrides, such as acetic anhydride, propyl anhydride, cyclohexanoic anhydride, benzoic anhydride, hippuric anhydride and the like. The preferred acylating agents are benzoyl chloride, benzene sulfonyl chloride, hippuryl chloride, acetyl chloride, cyclohexanoyl chloride, cyclopentanoyl chloride, and cycloheptanoyl chloride.

The amine groups can also be modified by the reaction of a carboxylic acid with coupling agents such as dicyclohexylcarbodiimide and the like. In a peptide one or more of the amino acids may be derivatized (an aldehyde or a ketone) and/or modified (acylated).


Also suitable as a carrier alone or in combination with the modified amino acid or peptide derivatives are the carbomethoxy modified amino acids carboxy-methyl-phenylalanine-leucine, 2-carboxy-3-phenylpropionyl-leucine, 2-benzylsuccinic acid and an actinonin. The actinonin compounds include actinonin or epiactinonin and derivatives thereof. These compounds have the formulas below:



15

Derivatives of these compounds are disclosed in U.S. Patent No. 5,206,384.

Actinonin derivatives have the formula:



wherein  $R^{11}$  is sulfoxymethyl or carboxyl or a substituted carboxy group selected from carboxamide, hydroxyaminocarbonyl and alkoxycarbonyl groups; and  $R^{12}$  is hydroxyl, alkoxy, hydroxyamino or sulfoxyamino group.

The modified amino acid derivatives or peptide derivatives can be readily prepared and modified by methods known to those skilled in the art. For example, the modified amino acid derivatives of the present invention may be prepared by reacting a single amino acid derivative or peptide derivative or mixtures of two or more amino acid or peptide derivatives, with an acylating agent or an amine modifying agent which reacts with free amino moieties present in the derivatives to form amides. The amino acid or peptide can be modified and subsequently derivatized, derivatized and subsequently modified, or simultaneously

modified and derivatized. Protecting groups may be used to avoid unwanted side reactions as would be known to those skilled in the art.

The modified amino acids and modified amino acid  
5 derivatives of the present invention may also be prepared by reacting single amino acids, mixtures of two or more kinds of amino acids, or amino acid esters with an amine modifying agent which reacts with free amino moieties present in the amino acids to form amides or sulfonamides. Amino acids and  
10 amino acid esters are readily available from a number of commercial sources such as Aldrich Chemical Co. (Milwaukee, WI, USA); Sigma Chemical Co. (St. Louis, MO, USA); and Fluka Chemical Corp. (Ronkonkoma, NY, USA).

For example, the amino acids can be dissolved in  
15 aqueous alkaline solution of a metal hydroxide, e.g., sodium or potassium hydroxide, and heated at a temperature ranging between about 5°C and about 70°C, preferably between about 10°C and about 40°C, for a period ranging between about 1 hour and about 4 hours, preferably about 2.5 hours. The  
20 amount of alkali employed per equivalent of  $\text{NH}_2$  groups in the amino acids generally ranges between about 1.25 and about 3 mmole, preferably between about 1.5 and about 2.25 mmole per equivalent of  $\text{NH}_2$ . The pH of the solution generally ranges between about 8 and about 13, preferably ranging between  
25 about 10 and about 12.

Thereafter, an amino modifying agent is added to the amino acid solution while stirring. The temperature of the mixture is maintained at a temperature generally ranging between about 5°C and about 70°C, preferably between about  
30 10°C and about 40°C, for a period ranging between about 1 and about 4 hours. The amount of amino modifying agent employed in relation to the quantity of amino acids is based on the moles of total free  $\text{NH}_2$  in the amino acids. In general, the amino modifying agent is employed in an amount  
35 ranging between about 0.5 and about 2.5 mole equivalents, preferably between about 0.75 and about 1.25 equivalents, per molar equivalent of total  $\text{NH}_2$  groups in the amino acids.



The reaction is quenched by adjusting the pH of the mixture with a suitable acid, e.g., concentrated hydrochloric acid, until the pH reaches between about 2 and about 3. The mixture separates on standing at room temperature to form a transparent upper layer and a white or off-white precipitate. The upper layer is discarded and modified amino acids are collected from the lower layer by filtration or decantation. The crude modified amino acids are then dissolved in water at a pH ranging between about 9 and about 13, preferably between about 11 and about 13. Insoluble materials are removed by filtration and the filtrate is dried in vacuo. The yield of modified amino acids generally ranges between about 30 and about 60%, and usually about 45%.

If desired, amino acid esters, such as, for example methyl or ethyl esters of amino acids, may be used to prepare the modified amino acids of the invention. The amino acid esters, dissolved in a suitable organic solvent such as dimethylformamide or pyridine, are reacted with the amino modifying agent at a temperature ranging between about 5°C and about 70°C, preferably about 25°C, for a period ranging between about 7 and about 24 hours. The amount of amino modifying agents used relative to the amino acid esters are the same as described above for amino acids.

Thereafter, the reaction solvent is removed under negative pressure and the ester functionality is removed by hydrolyzing the modified amino acid ester with a suitable alkaline solution, e.g. 1N sodium hydroxide, at a temperature ranging between about 50°C and about 80°C, preferably about 70°C, for a period of time sufficient to hydrolyze off the ester group and form the modified amino acid having a free carboxyl group. The hydrolysis mixture is then cooled to room temperature and acidified, e.g. aqueous 25% hydrochloric acid solution, to a pH ranging between about 2 and about 2.5. The modified amino acid precipitates out of solution and is recovered by conventional means such as filtration or decantation.

The modified amino acids may be purified by recrystallization or by fractionation on solid column supports. Suitable recrystallization solvent systems include acetonitrile, methanol and tetrahydrofuran.

5 Fractionation may be performed on a suitable solid column supports such as alumina, using methanol/n-propanol mixtures as the mobile phase; reverse phase column supports using trifluoroacetic acid/acetonitrile mixtures as the mobile phase; and ion exchange chromatography using water as the  
10 mobile phase. When anion exchange chromatography is performed, a subsequent 0-500 mM sodium chloride gradient is employed. The modified amino acids may also be purified by extraction with a lower alcohol such as methanol, butanol, or isopropanol to remove low molecular weight non-sphere  
15 making material.

Suitable modified amino acid derivatives include, but are not limited to, N-cyclohexanoyl-Phe aldehyde, N-acetyl-Phe-aldehyde, N-acetyl-Tyr ketone, N-acetyl-Lys ketone and N-acetyl-Leu ketone. Special mention is made of  
20 the modified amino acid derivative N-cyclohexanoyl phenylalanine aldehyde.

Special mention is made of compositions in which the biologically-active agent includes, calcitonin and the carrier includes acetyl phenylalanine aldehyde, carbomethoxy  
25 phenylalanylleucine and acetyl-Phe-Leu-Leu aldehyde.

Special mention is also made of a composition which includes 1.5  $\mu$ g/ml of the biologically-active agent calcitonin and the carrier includes 132 mg/ml of acetyl phenylalanine, 33 mg/ml of carbomethoxy phenylalanylleucine,  
30 and 25 mg/ml of acetyl-Phe-Leu-Leu-Arg aldehyde.

In one embodiment, the modified and/or modified derivatized amino acids may be used directly as a delivery carrier by simply mixing the carrier with the active ingredient prior to administration. In an alternative  
35 embodiment, the modified amino acids may be used to form microspheres containing the active agent. The modified and/or modified derivatized amino acids of the invention are particularly useful for the oral administration of certain

pharmacological agents, e.g., small peptide hormones, which, by themselves, do not pass or only pass slowly through the gastro-intestinal mucosa and/or are susceptible to chemical cleavage by acids and enzymes in the gastro-intestinal tract.

If the modified amino acids are to be converted into microspheres, the mixture is optionally heated to a temperature ranging between about 20 and about 50°C, preferably about 40°C, until the modified amino acid(s) dissolve. The final solution contains between from about 1 mg and about 2000 mg of modified amino acids per mL of solution, preferably between about 1 and about 500 mg per mL. The concentration of active agent in the final solution varies and is dependent on the required dosage for treatment. When necessary, the exact concentration can be determined by, for example, reverse phase HPLC analysis.

When the modified amino acids are used to prepare microspheres, another useful procedure is as follows: Modified amino acids are dissolved in deionized water at a concentration ranging between about 75 and about 200 mg/ml, preferably about 100 mg/ml at a temperature between about 25°C and about 60°C, preferably about 40°C. Particulate matter remaining in the solution may be removed by conventional means such as filtration.

Thereafter, the modified amino acid solution, maintained at a temperature of about 40°C, is mixed 1:1 (V/V) with an aqueous acid solution (also at about 40°C) having an acid concentration ranging between about 0.05 N and about 2 N, preferably about 1.7 N. The resulting mixture is further incubated at 40°C for a period of time effective for microsphere formation, as observed by light microscopy. In practicing this invention, the preferred order of addition is to add the modified amino acid solution to the aqueous acid solution.

Suitable acids for microsphere formation include any acid which does not

- (a) adversely effect the modified amino acids, e.g., initiate or propagate chemical decomposition;
- (b) interfere with microsphere formation;
- 5 (c) interfere with microsphere encapsulation of the cargo; and
- (d) adversely interact with the cargo.

Preferred acids for use in this invention include acetic acid, citric acid, hydrochloric acid, phosphoric  
10 acid, malic acid and maleic acid.

In practicing the invention, a microsphere stabilizing additive may be incorporated into the aqueous acid solution or into the amino acid solution prior to the microsphere formation process. With some drugs the presence  
15 of such additives promotes the stability and/or dispersibility of the microspheres in solution.

The stabilizing additives may be employed at a concentration ranging between about 0.1 and 5 % (w/v), preferably about 0.5 % (w/v). Suitable, but non-limiting,  
20 examples of microsphere stabilizing additives include gum acacia, gelatin, methyl cellulose, polyethylene glycol, and polylysine. The preferred stabilizing additives are gum acacia, gelatin and methyl cellulose.

Under the above conditions, the modified amino  
25 acid molecules form hollow or solid matrix type microspheres wherein the cargo is distributed in a carrier matrix or capsule type microspheres encapsulating liquid or solid cargo. If the modified amino acid microspheres are formed in the presence of a soluble material, e.g., a  
30 pharmaceutical agent in the aforementioned aqueous acid solution, this material will be encapsulated within the microspheres. In this way, one can encapsulate pharmacologically active materials such as peptides, proteins, and polysaccharides as well as charged organic  
35 molecules, e.g., antimicrobial agents, which normally have poor bioavailability by the oral route. The amount of pharmaceutical agent which may be encapsulated by the microsphere is dependent on a number of factors which

include the concentration of agent in the encapsulating solution, as well as the affinity of the cargo for the carrier.

The modified amino acid microspheres of the invention are pharmacologically harmless and do not alter the physiological and biological properties of the active agent. Furthermore, the encapsulation process does not alter the pharmacological properties of the active agent. Any pharmacological agent can be encapsulated within the amino acid microspheres. The system is particularly advantageous for delivering chemical or biological agents which otherwise would be destroyed or rendered less effective by conditions encountered within the body of the animal to which it is administered, before the microsphere reaches its target zone (*i.e.*, the area in which the contents of the microsphere are to be released) and pharmacological agents which are poorly absorbed in the gastro-intestinal tract. The target zones can vary depending upon the drug employed.

The particle size of the microsphere plays an important role in determining release of the active agent in the targeted area of the gastro-intestinal tract. The preferred microspheres have diameters between about  $\leq 0.1$  microns and about 10 microns, preferably between about 0.5 microns and about 5 microns. The microspheres are sufficiently small to release effectively the active agent at the targeted area within the gastro-intestinal tract. Small microspheres can also be administered parenterally by being suspended in an appropriate carrier fluid (*e.g.*, isotonic saline) and injected directly into the circulatory system, intramuscularly or subcutaneously. The mode of administration selected will vary, of course, depending upon the requirement of the active agent being administered. Large amino acid microspheres ( $>50$  microns) tend to be less effective as oral delivery systems.

The size of the microspheres formed by contacting modified amino acid with water or an aqueous solution containing active agents can be controlled by manipulating a

variety of physical or chemical parameters, such as the pH, osmolarity or ionic strength of the encapsulating solution, size of the ions in solution and by the choice of acid used in the encapsulating process.

5 Typically, the pharmacological compositions of the present invention are prepared by mixing an aqueous solution of the carrier with an aqueous solution of the active ingredient, just prior to administration. Alternatively, the carrier and biologically active ingredient can be  
10 admixed during the manufacturing process. The solutions may optionally contain additives such as phosphate buffer salts, citric acid, acetic acid, gelatin and gum acacia.

In practicing the invention, stabilizing additives may be incorporated into the carrier solution. With some  
15 drugs, the presence of such additives promotes the stability and dispersibility of the agent in solution.

The stabilizing additives may be employed at a concentration ranging between about 0.1 and 5 % (W/V), preferably about 0.5 % (W/V). Suitable, but non-limiting,  
20 examples of stabilizing additives include gum acacia, gelatin, methyl cellulose, polyethylene glycol, and polylysine. The preferred stabilizing additives are gum acacia, gelatin and methyl cellulose.

The amount of active agent in the composition  
25 typically is a pharmacologically or biologically effective amount. However, the amount can be less than a pharmacologically or biologically effective amount when the composition is used in a dosage unit form, such as a capsule, a tablet or a liquid, because the dosage unit form  
30 may contain a multiplicity of carrier/biologically-active agent compositions or may contain a divided pharmacologically or biologically effective amount. The total effective amounts will be administered by cumulative units containing in total pharmacologically or biologically  
35 active amounts of biologically-active agent.

The total amount of biologically-active agent to be used can be determined by those skilled in the art. However, it has surprisingly been found that with certain

biologically-active agents, such as calcitonin, the use of the presently disclosed carriers provides extremely efficient delivery. Therefore, lower amounts of biologically-active agent than those used in prior dosage unit forms or delivery systems can be administered to the subject, while still achieving the same blood levels and therapeutic effects.

The amount of carrier in the present composition is a delivery effective amount and can be determined for any particular carrier or biologically-active agent by methods known to those skilled in the art.

Dosage unit forms can also include any of excipients; diluents; disintegrants; lubricants; plasticizers; colorants; and dosing vehicles, including, but not limited to water, 1,2-propane diol, ethanol, olive oil, or any combination thereof.

Administration of the present compositions or dosage unit forms is oral or by intraduodenal injection.

## 20 EXAMPLES

The invention will now be illustrated in the following non-limiting examples which are illustrative of the invention but are not intended to limit the scope of the invention.

25

### EXAMPLE 1

#### PREPARATION OF N-CYCLOHEXANOYLPHENYLALANINE ALDEHYDE:

Phenylalanine methyl ester (1 g., 0.0046 moles) was dissolved in pyridine 5 mL. Cyclohexanoyl chloride (0.62 mL) was added and the mixture was stirred for 2 hours. The reaction mixture was poured onto hydrochloric acid (1N) and crushed ice. The aqueous mixture was extracted twice with toluene. The combined toluene extracts were concentrated in vacuo to give 1.1 g of crude N-cyclohexanoylphenylalanine methyl ester.

N-Cyclohexanoylphenylalanine methyl ester (0.5 g) was dissolved in ethylene glycol dimethyl ether (20 mL). The solution was cooled to -70°C and diisobutylaluminum

hydride (2.04 mL of a 1.5M solution in toluene) was added. The resulting reaction mixture was stirred at -70°C for 2 hours. The reaction was quenched by dropwise addition of 2N hydrochloric acid. The mixture was extracted with cold ethyl acetate. The ethyl acetate solution was washed with brine and dried over sodium sulfate. Concentration in vacuo furnished a white solid which was purified by silica gel chromatography. <sup>1</sup>H NMR(300 MHz, DMSO-d<sub>6</sub>): 9.5 (s, 1H), 8.2 (d, 1H), 7.2 (m, 5H), 4.2 (m, 1H), 3.2 (d, 1H), 2.7 (d, 1H), 2.1 (m, 1H), 1.6 (br. m, 4H), 1.2 (br. m, 6H). IR (KBr): 3300, 3050, 2900, 2850, 2800, 1700, 1600, 1500 cm<sup>-1</sup>. Mass Spec.: M+1 m/e 261.

## 15 EXAMPLE 2

### PREPARATION OF N-ACETYLPHENYLALANINE ALDEHYDE:

N-Acetylphenylalanine methyl ester (4.2 g, 19 mmol) was dissolved in ethylene glycol dimethyl ether. The solution was cooled to -70°C and diisobutylaluminum hydride (25.3 mL of a 1.5M solution in toluene, 39 mmol) was added. The resulting reaction mixture was stirred at -70°C for 2 hours. The reaction was quenched by addition of 2N hydrochloric acid. The mixture was extracted 4 times with cold ethyl acetate and 4 times with toluene. The extracts were combined, washed with brine and dried over magnesium sulfate. Concentration in vacuo followed by silica gel chromatography furnished 2.7 g of a white solid. The NMR was identical to that reported in the literature, *Biochemistry*, 1979, 18, 921-926.

30

## EXAMPLE 3

### PREPARATION OF 3-ACETAMIDO-4-(p-HYDROXY)PHENYL-2-BUTANONE (N-ACETYLTYROSINONE):

A mixture of tyrosine (28.9 g, 16 mmol), acetic anhydride (97.9 g, 96 mmol) and pyridine (35g, 16 mmol) were heated to 100 °C for 1 hour. The reaction mixture was concentrated in vacuo to furnish a yellow oil. The oil was distilled at reduced pressure to furnish 29.9 g or an oil.



<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): NMR (d<sub>6</sub>-DMSO); 8.2 (d, 1H), 7.3 (d, 2H), 7.0 (d, 2H), 4.4 (m, 1H), 3.1 (dd, 1H), 2.7 (dd, 1H), 2.3 (s, 3H), 1.8 (s, 3H).

5 EXAMPLE 4

PREPARATION OF 3-ACETAMIDO-7-AMINO-2-BUTANONE  
(N-ACETYLLYSINONE):

Following the procedure of Example 3 lysine was converted to N-acetyllysine.

10 <sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 8.1 (d, 1H), 7.8 (br.m. 1H), 4.1 (m, 1H), 3.0 (m, 2H), 2.0 (s, 3H), 1.9 (s, 3H) and 1.3 (br.m, 6H).

EXAMPLE 5

PREPARATION OF 3-ACETAMIDO-5-METHYL-2-BUTANONE  
(N-ACETYLLLEUCINONE):

Following the procedure of Example 3 leucine was converted to N-acetylleucine.

<sup>1</sup>H NMR (DMSO-d<sub>6</sub>): 8.1 (d, 1H), 4.2 (m, 1H), 2.0 (s, 3H), 1.8 (s, 3H), 0.8 (d, 6H).

20

EXAMPLE 6

MODIFICATION OF 4-(4-AMINOPHENYL)BUTYRIC ACID USING BENZENE  
SULFONYL CHLORIDE

4-(4-Aminophenyl)butyric acid, (20 g 0.11 moles)  
25 was dissolved in 110 mL of aqueous 2N sodium hydroxide solution. After stirring for about 5 minutes at room temperature, benzene sulfonyl chloride (14.2 mL, 0.11 moles) was added dropwise into the amino acid solution over a 15 minute period. After stirring for about 3 hours at room  
30 temperature the mixture was acidified to pH 2 by addition of hydrochloric acid. This furnished a light brown precipitate which was isolated by filtration. The precipitate was washed with warm water and dried. The yield of 4-(phenyl-sulfonamido)4-phenylbutyric acid was 24.3 g (69%). The  
35 melting point was 123-25°C.

If necessary, the modified amino acids can be purified by recrystallization and/or chromatography.

EXAMPLE 7MODIFICATION OF 4-AMINOBENZOIC ACID USING BENZENE SULFONYL CHLORIDE

Following the procedure of Example 6 4-aminobenzoic acid was converted to 4-(phenylsulfonamido)benzoic acid.

EXAMPLE 8

10 MODIFICATION OF 4-AMINOPHENYLACETIC ACID, 4-AMINOHIPPURIC ACID, AND 4-AMINOMETHYLBENZOIC ACID USING BENZENE SULFONYL CHLORIDE

Following the procedure of Example 6, 4-aminophenylacetic acid, 4-aminohippuric acid, and 4-aminomethylbenzoic acid were converted to 4-(phenylsulfonamido)-phenylacetic acid, 4-(phenylsulfonamido)hippuric acid, and 4-(phenylsulfonamidomethyl)benzoic acid respectively.

EXAMPLE 9MODIFICATION OF AMINO ACIDS WITH BENZENE SULFONYL CHLORIDE

20 A mixture of sixteen amino acids were prepared prior to chemical modification. The constituents of the mixture are summarized in Table 1. 65 grams of the amino acid mixture (total concentration of [-NH<sub>2</sub>] groups = 0.61 moles) was dissolved in 760 mL of 1N sodium hydroxide solution (0.7625 equivalents) at room temperature. After 25 stirring for 20 minutes, benzene sulfonyl chloride (78 ml, 1 eq.) was added over a 20 minute period. The reaction mixture was then stirred for 2.5 hours, without heating. As some precipitation had occurred, additional NaOH solution 30 (2N) was added to the solution until it reached pH 9.3. The reaction mixture stirred overnight at room temperature. Thereafter, the mixture was acidified using dilute hydrochloric acid (38%, 1:4) and a cream colored material precipitated out. The resulting precipitate was isolated by 35 decantation and dissolved in sodium hydroxide (2N). This solution was then reduced in vacuo to give a yellow solid, which was dried on the lyophilizer.

**TABLE 1: Amino Acid Composition**

5	Amino Acid	Weight (g)	% of Total Weight	No. of moles of each Amino Acid ( $\times 10^{-2}$ )	No. of Moles of - $[-NH_2]$
	Thr	2.47	3.8	2.07	2.07
	Ser	2.25	3.46	2.1	2.1
	Ala	4.61	7.1	5.17	5.17
10	Val	4.39	6.76	3.75	3.75
	Met	0.53	0.82	0.35	0.35
	Ile	2.47	3.8	0.35	0.36
	Leu	3.86	5.94	2.95	2.95
	Tyr	1.03	1.58	0.56	0.56
15	Phe	4.39	6.76	0.27	0.27
	His	2.47	3.8	1.6	3.2
	Lys	4.94	7.6	3.4	6.8
	Arg	5.13	7.9	2.95	5.90
	Glutamine	9.87	15.18	6.76	13.42
20	Glutamic Acid	9.87	15.18	6.70	6.70
	Asparagine	3.32	5.11	2.51	5.02
25	Aspartic Acid	3.32	5.11	2.50	2.50

**EXAMPLE 10****MODIFICATION OF A MIXTURE OF FIVE AMINO ACIDS USING BENZENE SULFONYL CHLORIDE**

An 86.1g (0.85 moles of  $NH_2$ ) mixture of amino acids (see Table 2) was dissolved in 643 mL (1.5 eq.) of aqueous 2N sodium hydroxide solution. After stirring for 30 minutes at room temperature, benzene sulfonyl chloride (108 mL, 0.86 moles) was added portionwise into the amino acid solution over a 15 minute period. After stirring for 2.5 hours at room temperature, the pH of the reaction mixture (pH 5) was adjusted to pH 9 with additional 2N sodium hydroxide solution. The reaction mixture stirred overnight at room temperature. Thereafter, the pH of the reaction mixture was

adjusted to pH 2.5 by addition of dilute aqueous hydrochloric acid solution (4:1, H<sub>2</sub>O:HCl) and a precipitate of modified amino acids formed. The upper layer was discarded and the resulting yellow precipitate was isolated by decantation, washed with water and dissolved in 2N sodium hydroxide (2N). The solution was reduced in vacuo to give a yellow solid which was lyophilized overnight. The yield of crude modified amino acid was 137.9 g.

10

Table 2

15

Amino Acid	Moles of Amino Acid (x 10 <sup>2</sup> )	Moles of [-NH <sub>2</sub> ] <sup>+</sup> x10 <sup>2</sup>
Valine	7.5	7.5
Leucine	10.7	10.5
Phenylalanine	13.4	13.4
Lysine	21.0	42.0
Arginine	6.0	12.0

EXAMPLE 11MODIFICATION OF A MIXTURE OF FIVE AMINO ACIDS USING BENZOYL CHLORIDE

An 86 g (0.85 moles of  $\text{NH}_2$ ) mixture of amino acids (see Table 2 in Example 10) was dissolved in 637 mL (1.5 eq.) of aqueous 2N sodium hydroxide solution. After stirring for 10 minutes at room temperature, benzoyl chloride (99 mL, 0.85 moles) was added portionwise into the amino acid solution over a 10 minute period. After stirring for 2.5 hours at room temperature, the pH of the reaction mixture (pH 12) was adjusted to pH 2.5 using dilute hydrochloric acid (4:1,  $\text{H}_2\text{O}:\text{HCl}$ ) and a precipitate of modified amino acids formed. After settling for 1 hour, the resulting precipitate was isolated by decantation, washed with water and dissolved in sodium hydroxide (2N). This solution was then reduced in vacuo to give crude modified amino acids as a white solid (220.5 g).

EXAMPLE 12MODIFICATION OF L-VALINE USING BENZENE SULFONYL CHLORIDE

L-Valine (50 g, 0.43 mol) was dissolved in 376 mL (0.75 eq.) of aqueous 2N sodium hydroxide by stirring at room temperature for 10 minutes. Benzene sulfonyl chloride (68.7 mL, 0.38 mol, 1.25 eq.) was then added to the amino acid solution over a 20 minute period at room temperature. After stirring for 2 hours at room temperature, a precipitate appeared. The precipitate was dissolved by adding 200 mL of additional 2N sodium hydroxide solution. After stirring for an additional 30 minutes, dilute aqueous hydrochloric acid solution (4:1,  $\text{H}_2\text{O}:\text{HCl}$ ) was added until the pH of the reaction mixture reached 2.6. A precipitate of modified amino acids formed and was recovered by decantation. This material was dissolved in 2N sodium hydroxide and dried in vacuo to give a white solid. Yield of crude modified amino acids = 84.6 g, 77%).

EXAMPLE 13MODIFICATION OF PHENYLALANINE METHYL ESTER USING HIPPURYL CHLORIDE

L-Phenylalanine Methyl Ester Hydrochloride (15 g, 0.084 mole) was dissolved in dimethylformamide (DMF) (100 mL) and to this was added pyridine (30 mL). A solution of hippuryl chloride (16.6 g, 0.084 moles in 100 mL DMF) was immediately added to the amino acid ester solution in two portions. The reaction mixture was stirred at room temperature overnight. The reaction mixture was then reduced in vacuo and dissolved in 1N aqueous sodium hydroxide. The solution was heated at 70°C for 3 hours in order to hydrolyze the methyl ester to a free carboxyl group. Thereafter, the solution was acidified to pH 2.25 using dilute aqueous hydrochloric acid solution (1:3 HCl/H<sub>2</sub>O). A gum-like precipitate formed and this was recovered and dissolved in 1N sodium hydroxide. The solution was reduced in vacuo to afford 18.6 g of crude modified amino acid product (Yield 18.6 g). After recrystallization from acetonitrile, pure modified phenylalanine (12 g) was recovered as a white powder. m.p. 223-225°C.

EXAMPLE 14PREPARATION OF DOSING SOLUTIONS:

In a test tube 568 mg of acetyl phenylalanine aldehyde, 132 mg of carbomethoxy phenylalanylleucine and 100 mg acetyl-Phe-Leu-Leu-Arg aldehyde were added to 2.9 mL of 15% ethanol. The solution was stirred and NaOH (1.0 N) was added to raise the pH to 7.2. Water was added to bring the total volume to 4.0 mL. The sample had a carrier concentration of 200 mg/mL. Calcitonin (6 µg) was added to the solution. The total calcitonin concentration was 1.5 µg/mL.

Following a similar procedure a second solution having 668 mg of acetyl phenylalanine aldehyde and 132 mg of carbomethoxy phenylalanylleucine as the carrier composition and a third solution having as the carrier acetyl phenyl-

alanine aldehyde. Each solution had a calcitonin concentration of 1.5  $\mu\text{g/mL}$ .

EXAMPLE 15

5 PREPARATION OF MODIFIED AMINO ACID/SALMON CALCITONIN  
COMPOSITIONS

(a) Preparation of Modified Amino acid microspheres  
containing encapsulated Salmon Calcitonin

10 The modified amino acid mixture, prepared in accordance with Example 9, was dissolved at 40°C in distilled water (pH 7.2) at a concentration of 100 mg/ml. The solution was then filtered with a 0.2 micron filter and the temperature was maintained at 40°C. Salmon calcitonin  
15 (Sandoz Corp., Basil, Switzerland) was dissolved in an aqueous solution of citric acid (1.7N) and gelatin (5%) at a concentration of 150 mg/ml. This solution was then heated to 40°C. The two heated solutions were then mixed 1:1 (v/v). The resulting microsphere suspension was then  
20 filtered with glass wool and centrifuged for 50 minutes at 1000 g. The pellet was resuspended with 0.85N citric acid to a volume 5 to 7 fold less than the original volume. Salmon calcitonin concentration of the resuspended pellet was determined by HPLC. Additional microspheres were made  
25 according to the above procedure without salmon calcitonin. These "empty microspheres" were used to dilute the encapsulated salmon calcitonin microsphere preparation to a final dosing suspension for animal testing.

30 (b) Preparation of a Soluble Modified Amino acid  
carrier/Salmon Calcitonin system

A soluble amino acid dosing preparation containing salmon calcitonin was prepared by dissolving the modified amino acid material in distilled water (pH 8) to an appropriate concentration. The solution was heated to 40°C  
35 and then filtered with a 0.2 micron filter. Salmon calcitonin, also dissolved in distilled water, was then added to the modified amino acid solution prior to oral administration.

EXAMPLE 16IN VIVO EXPERIMENTS IN RATS

For each sample six fasted rats were anesthetized. The rats were administered, by oral gavage, one of the  
5 calcitonin/carrier dosages prepared in Example 15. The calcitonin concentration in each sample was 1.5 µg/ml. Each rat was administered a dosage of two (2) mL/kg each. Blood samples were collected serially from the tail artery. Serum calcium was determined by testing with a Demand™ Calcium Kit  
10 (available from Sigma Chemical Company, St. Louis, Missouri, USA). The results of the test are illustrated in Figure 1.

EXAMPLE 17

Three samples having 400 mg/kg of acetyl-Leu  
15 aldehyde and 10 µg/kg of calcitonin, 400 mg/kg of acetyl-Phe aldehyde and 10 µg/kg of calcitonin, 200 mg/kg of acetyl-Leu aldehyde, 200 mg/kg of acetyl-Phe aldehyde and 10 µg/kg of calcitonin, respectively were prepared. The samples were given to fasted rats as described in Example 16. The  
20 results of the test are illustrated graphically in Figure 2.

EXAMPLE 18

Two samples having 350 mg/kg of acetyl-Phe aldehyde, 50 mg/kg of carbomethoxy-Phe-Leu-OH and 3 µg/kg of  
25 calcitonin, 400 mg/kg of acetyl-Phe aldehyde, 50 mg/kg of carbomethoxy-Phe-Leu-OH and 10 µg/kg of calcitonin, respectively were prepared. The samples were given to fasted rats as described in Example 16. The results of the test are illustrated in Figure 3.

30

EXAMPLE 19

Three samples having 284 mg/kg of acetyl-Phe aldehyde and 66 mg/kg acetyl-Leu-Leu-Arg aldehyde, 50 mg/kg of carbomethoxy-Phe-Leu-OH and 3 µg/kg of calcitonin in  
35 propylene glycol, 284 mg/kg of acetyl-Phe aldehyde and 66 mg/kg acetyl-Leu-Leu-Arg aldehyde, 50 mg/kg of carbomethoxy-Phe-Leu-OH and 3 µg/kg of calcitonin and 3 µg/kg of calcitonin, in aqueous ethanol, respectively were prepared.



The samples were given to fasted rats as described in Example 16. The results of the test are illustrated in Figure 4.

5 EXAMPLE 20

Three samples having 400 mg/kg of 4-(phenylsulfon-amido)-4-phenylbutyric acid and 1.5  $\mu$ g/kg of calcitonin in propylene glycol, 200 mg/kg of 4-(phenylsulfonamido)-4-phenylbutyric acid, 200 mg/kg of acetyl-Phe aldehyde and  
10 1.5  $\mu$ g/kg of calcitonin in aqueous ethanol, respectively were prepared. The samples were given to fasted rats by intraduodenal injection. The results of the test are illustrated in Figure 5.

15 EXAMPLE 21

Samples having 600 mg/kg of acetyl-Phe aldehyde and 10  $\mu$ g/kg of calcitonin in aqueous ethanol, and 3  $\mu$ g/kg of calcitonin, 200 mg/kg of acetyl-Phe aldehyde, 200 mg/kg N-acetyllysine, 200 mg/kg acetyl-Leu aldehyde and 10  $\mu$ g/kg  
20 of calcitonin were prepared. The samples were given to fasted rats as described in Example 16. The results of the test are illustrated in Figure 6.

EXAMPLE 22

25 Three samples having 200 mg/kg of acetyl-Phe aldehyde and 3  $\mu$ g/kg of calcitonin, in aqueous ethanol, dimethyl sulfoxide (DMSO), and olive oil, respectively, were prepared. Additionally a sample of 3  $\mu$ g/kg of calcitonin in DMSO alone was prepared. The samples were given to rats by  
30 intraduodenal injection. The results of the test are illustrated in Figure 7.

EXAMPLE 23

A sample having 400 mg/kg of cyclohexanoyl-Phe  
35 aldehyde and 3  $\mu$ g/kg of calcitonin in aqueous ethanol was prepared. The sample was given to fasted rats as described in Example 16. The results of the test are illustrated in Figure 8.

EXAMPLE 24

In vivo evaluation of modified amino acid microspheres containing encapsulated calcitonin and soluble modified amino acid carrier/calcitonin system, prepared as described in Example 16, were evaluated in rats. Rats were gavaged with the oral dosing preparations and blood samples were withdrawn at various time intervals for serum calcium concentration determinations.

Nine rats are divided into three groups as follows:

1. calcitonin microspheres: 10 ug calcitonin/kg body weight by oral gavage (3 rats);
2. calcitonin microspheres: 30 ug calcitonin/kg body weight by oral gavage (3 rats); and
3. soluble modified amino acid/calcitonin system: 30 ug calcitonin/kg body weight by oral gavage (3 rats). The rats were pre-dosed with 0.7 meq of aqueous sodium bicarbonate solution prior to administration of the soluble system.

Oral gavage dosing of rats is performed. Calcitonin microspheres are prepared immediately prior to dosing and Group 1 rats and Group 2 rats each receive an appropriate dosage of the microsphere suspension. Group 3 rats receives the unencapsulated calcitonin/modified amino acid system. Approximately 0.5 ml of blood is withdrawn from each rat just prior to dosing ("0" time) and 1 h, 2 h and 3 h post-dosing. Serum from the blood samples are stored at -20°C.

The calcium levels of thawed serum taken from group 1-3 rats are analyzed by conventional methods. Experimental results in rats have demonstrated a significant increase in pharmacological activity (i.e., decreasing serum calcium levels) when calcitonin is orally administered either as a encapsulate in modified amino acid microspheres or a mixture with modified amino acids as compared to basal levels. As shown in Figure 9, soluble modified amino acid solution containing salmon calcitonin demonstrated a significant increase in pharmacological activity (i.e.,

decreasing serum calcium levels) when compared to basal levels after oral administration.

#### EXAMPLE 25

5 Two samples having 366 mg/kg of acetyl-Phe aldehyde, 33 mg/kg of actinonin and 10 µg/kg of calcitonin, 366 mg of acetyl-Phe aldehyde, 33 mg/kg of carbomethoxy-Phe-Leu-OH and 10 µg/kg of calcitonin, respectively, were prepared. The samples were given to fasted rats as  
10 described in Example 14. The results of the test are illustrated in Figure 10.

#### EXAMPLE 26

Two samples having 400 mg/kg of 4-  
15 (phenylsulfonamido)-4-phenylbutyric acid and 3 µg/kg of calcitonin, 400 mg/kg of 4-(phenylsulfonamido)-4-phenylbutyric acid and 10 µg/kg of calcitonin, respectively were prepared. The samples were given to fasted rats as described in Example 14. The results of the test are  
20 illustrated in Figure 11.

#### EXAMPLE 27

Two samples having 400 mg/kg of 3-  
(phenylsulfonamido)benzoic acid and 10 µg/kg of calcitonin,  
25 400 mg/kg of 4-(phenylsulfonamido)hippuric acid and 10 µg/kg of calcitonin, respectively were prepared. The samples were given to fasted rats as described in Example 14. The results of the test are illustrated in Figure 12.

30

#### EXAMPLE 28

Two samples having 400 mg/kg of 4-  
(phenylsulfonamido)-4-phenylbutyric acid and 10 µg/kg of calcitonin, 400 mg/kg of 4-(phenylsulfonamido)benzoic acid  
35 and 10 µg/kg of calcitonin, respectively were prepared. The samples were given to fasted rats as described in Example 14. The results of the test are illustrated in Figure 13.

EXAMPLE 29

Two samples having 400 mg/kg of 4-(phenylsulfonamido)-4-phenylbutyric acid and 10 µg/kg of calcitonin, 400 mg/kg of 4-(phenylsulfonamido)phenylacetic acid and 10 µg/kg of calcitonin, respectively were prepared. The samples were given to fasted rats as described in Example 14. The results of the test are illustrated in Figure 14.

In Vivo EVALUATION OF INTERFERON PREPARATIONS IN RATS

Following the procedure described herein samples containing the carriers of the subject invention, in a Trizma® hydrochloride buffer solution (Tris-HCl) at a pH of about 7-8, and interferon α2b were prepared. The animals were administered the drug by oral gavage. The delivery was evaluated by using an ELISA assay for human interferon α.

EXAMPLE 30

A sample having 800 mg/kg of 4-(phenylsulfonamido)-4-phenylbutyric acid in a buffered solution and 1000 µg/kg of interferon α2b was prepared. The sample was given to fasted rats as described in Example 14. The results of the test are illustrated in Figure 15.

EXAMPLE 31

A sample having 400 mg/kg of 4-(phenylsulfonamidomethyl)benzoic acid in a buffered solution and 1000 µg/kg of interferon α2b was prepared. The sample was given to fasted rats as described in Example 14. The results of the test are illustrated in Figure 16.

EXAMPLE 32

A sample having 800 mg/kg of 4-(phenylsulfonamido)phenylacetic acid in a buffered solution and 1000 µg/kg of interferon α2b was prepared. The sample was given to fasted rats as described in Example 14. The results of the test are illustrated in Figure 17.

EXAMPLE 33

A sample having 600 mg/kg of 4-(phenylsulfonamido)hippuric acid in a buffered solution and 1000 µg/kg of interferon α2b was prepared. The sample was given to fasted rats as described in Example 14. The results of the test  
5 are illustrated in Figure 18.

#### In Vivo EVALUATION OF GROWTH HORMONE PREPARATIONS IN RATS

Following the procedure described herein samples containing the carriers of the subject invention and growth  
10 hormone were prepared. The animals were administered the drug by oral gavage. The delivery was evaluated by using an ELISA assay for growth hormone.

#### EXAMPLE 34

A sample having 1000 mg/kg of 4-(phenylsulfonamido)-4-phenylbutyric acid and 1 mg/kg of growth hormone was prepared. The sample was given to hypophysectomized rats as  
15 described in Example 14. The results of the test are illustrated in Figure 19.

20

#### EXAMPLE 35

A sample having 500 mg/kg of 4-(phenylsulfonamido)-4-phenylbutyric acid and 1 mg/kg of growth hormone was prepared. In a comparison a group of hypophysectomized rats  
25 were given samples of growth hormone without a carrier. The samples were given to hypophysectomized rats as described in Example 14. The results of the test are illustrated in Figure 20.

#### EXAMPLE 36

Two samples having 500 mg/kg of 4-(phenylsulfonamido)-4-phenylbutyric acid and 6 mg/kg of growth hormone were prepared. The samples were given to normal rats as  
30 described in Example 14. The results of the tests are illustrated in Figure 21.

35

In Vivo EVALUATION OF CROMOGLYCOLATE PREPARATIONS IN RATSExample 37

Following the procedure described herein samples containing the carriers of the subject invention and disodium cromoglycolate were prepared. The sample, in 0.85N citric acid and 0.5% acacia, contained 200 mg/kg of 4-(phenylsulfonamido)-4-phenylbutyric acid and 50 mg/kg of disodium cromoglycate. The animals were administered the drug by oral gavage. The delivery was evaluated by using the procedure described by A. Yoshimi in *Pharmacobio-Dyn.*, 15, pages 681-686, (1992). The results of the tests are illustrated in Figure 22.

As clearly illustrated by the data in the Examples and Figures the use of compositions of the subject invention show significant advantages for the delivery of biologically active agents.

All patents, applications, and publications mentioned herein are incorporated by reference herein.

Many variations of the present invention will suggest themselves to those skilled in the art in light of the above detailed disclosure. For example, poly (amino acids) which are formed by a bond other than an amide bond, e.g., an ester or an anhydride linkage, may be derivatized and modified for use as carriers in accordance with the present invention. All such modifications are within the full intended scope of the appended claims.

WHAT IS CLAIMED IS:

1. A composition comprising:
  - (A) at least one biologically-active agent; and
  - (B) at least one carrier comprising
    - (a) (i) at least one acylated aldehyde of an amino acid,
    - (ii) at least one acylated ketone of an amino acid,
    - (iii) at least one acylated aldehyde of a peptide,
    - (iv) at least one acylated ketone of a peptide, or
    - (v) any combination of (a) (i), (a) (ii), (a) (iii) and (a) (iv);
  - (b) (i) carboxymethyl-phenylalanine-leucine,
  - (ii) 2-carboxy-3-phenylpropionyl-leucine,
  - (iii) 2-benzylsuccinic acid, or
  - (iv) an actinonin, or
  - (v) a compound having the formula:
 
$$\text{Ar}-\text{Y}-(\text{R}^1)_n-\text{OH}$$

wherein: Ar is a substituted or unsubstituted phenyl or naphthyl;

Y is  $\text{—}\overset{\text{O}}{\parallel}\text{C—}$  or  $\text{—SO}_2\text{—}$ ;

$\text{R}^1$  is  $\text{—N}(\text{R}^4)\text{—R}^3\text{—}\overset{\text{O}}{\parallel}\text{C—}$ ; wherein:

$\text{R}^3$  is  $\text{C}_1$  to  $\text{C}_{24}$  alkyl,  $\text{C}_1$  to  $\text{C}_{24}$  alkenyl, phenyl, naphthyl, ( $\text{C}_1$  to  $\text{C}_{10}$  alkyl) phenyl, ( $\text{C}_1$  to  $\text{C}_{10}$  alkenyl) phenyl, ( $\text{C}_1$  to  $\text{C}_{10}$  alkyl) naphthyl, ( $\text{C}_1$  to  $\text{C}_{10}$  alkenyl) naphthyl, phenyl ( $\text{C}_1$  to  $\text{C}_{10}$  alkyl), phenyl ( $\text{C}_1$  to  $\text{C}_{10}$  alkenyl), naphthyl ( $\text{C}_1$  to  $\text{C}_{10}$  alkyl) and naphthyl ( $\text{C}_1$  to  $\text{C}_{10}$  alkenyl);

$\text{R}^3$  is optionally substituted with  $\text{C}_1$  to  $\text{C}_4$  alkyl,  $\text{C}_1$  to  $\text{C}_4$  alkenyl,  $\text{C}_1$  to  $\text{C}_4$  alkoxy,  $\text{—OH}$ ,  $\text{—SH}$ ,  $\text{—CO}_2\text{R}^5$ , cycloalkyl, cycloalkenyl, heterocyclic, aryl, alkaryl, heteroaryl or heteroalkaryl or any combination thereof;

$\text{R}^5$  is hydrogen,  $\text{C}_1$  to  $\text{C}_4$  alkyl or  $\text{C}_1$  to  $\text{C}_4$  alkenyl;

36 R<sup>3</sup> is optionally interrupted by oxygen,  
37 nitrogen, sulfur or any combination thereof; and  
38 R<sup>4</sup> is hydrogen, C<sub>1</sub> to C<sub>4</sub> alkyl or C<sub>1</sub> to C<sub>4</sub>  
39 alkenyl;  
40 and n is from 1 to 5;  
41 (vi) any combination of (b) (i), (b) (ii),  
42 (b) (iii), (b) (iv) and (b) (v); or  
43 (c) a combination of (a) and (b).

1 2. The composition according to claim 1 wherein R<sup>3</sup> is  
2 substituted with a substituent selected from the group  
3 consisting of C<sub>1</sub> to C<sub>4</sub> alkyl, C<sub>1</sub> to C<sub>4</sub> alkenyl, C<sub>1</sub> to C<sub>4</sub>  
4 alkoxy, -OH, -SH, -CO<sub>2</sub>R<sup>5</sup> or any combination thereof.

1 3. The composition according to claim 1, wherein said  
2 biologically-active agent is selected from the group  
3 consisting of a peptide, a mucopolysaccharide, a  
4 carbohydrate, a lipid, a pesticide or any combination  
5 thereof.

1 4. The composition according to claim 1, wherein said  
2 biologically-active agent is selected from the group  
3 consisting of human growth hormone, bovine growth hormone,  
4 growth hormone-releasing hormone, an interferon,  
5 interleukin-II, insulin, heparin, calcitonin,  
6 erythropoietin, atrial natriuretic factor, an antigen, a  
7 monoclonal antibody, somatostatin, adrenocorticotropin,  
8 gonadotropin releasing hormone, oxytocin, vasopressin,  
9 cromolyn sodium, vancomycin, desferrioxamine (DFO), or any  
10 combination thereof.

1 5. The composition according to claim 4, wherein said  
2 biologically-active agent comprises an interferon,  
3 interleukin-II, insulin, heparin, calcitonin, oxytocin,  
4 vasopressin, cromolyn sodium, vancomycin, DFO or any  
5 combination thereof.



1           6. The composition according to claim 5, wherein said  
2 biologically-active agent is calcitonin.

1           7. The composition according to claim 1, wherein said  
2 amino acid is a naturally occurring amino acid.

1           8. The composition according to claim 1, wherein said  
2 amino acid is a synthetic amino acid.

1           9. The composition according to claim 1, wherein said  
2 amino acid is an  $\alpha$ -amino acid.

1           10. The composition according to claim 1, wherein said  
2 amino acid is a non- $\alpha$ -amino acid.

1           11. The composition according to claim 1, wherein said  
2 amino acid is selected from the group consisting of alanine,  
3 arginine, asparagine, aspartic acid, citrulline, cysteine,  
4 cystine, glutamine, glycine, histidine, isoleucine, leucine,  
5 lysine, methionine, ornithine, phenylalanine, phenylglycine,  
6 proline, serine, threonine, tryptophan tyrosine, valine,  
7 hydroxyproline,  $\gamma$ -carboxyglutamate, O-phosphoserine,  $\beta$ -  
8 alanine,  $\alpha$ -aminobutyric acid,  $\gamma$ -aminobutyric acid,  $\alpha$ -  
9 aminoisobutyric acid, 4-(4-aminophenyl)butyric acid, (amino-  
10 phenyl)acetic acid, aminobenzoic acid, 4-aminohippuric acid,  
11 (aminomethyl)benzoic acid  $\epsilon$ -aminocaproic acid, 7-  
12 aminoheptanoic acid,  $\beta$ -aspartic acid,  $\gamma$ -glutamic acid,  
13 cysteine (ACM),  $\epsilon$ -lysine,  $\epsilon$ -lysine (A-Fmoc), methionine  
14 sulfone, norleucine, norvaline, ornithine, d-ornithine,  
15 p-nitro-phenylalanine, hydroxy proline, and thioproline.

1           12. The composition according to claim 11, wherein said  
2 amino acid is selected from the group consisting of  
3 arginine, leucine, lysine, phenylalanine, tyrosine, valine,  
4 phenylglycine, 4-(4-aminophenyl)butyric acid, 4-(4-amino-  
5 phenyl)acetic acid and aminobenzoic acid.

1        13. The composition according to claim 1, wherein said  
2 peptide is selected from the group consisting of a  
3 di-peptide, a tri-peptide, a tetra-peptide, or a  
4 penta-peptide.

1        14. The composition according to claim 1, wherein said  
2 peptide comprises at least one naturally occurring amino  
3 acid.

1        15. The composition according to claim 1, wherein said  
2 peptide comprises at least one synthetic amino acid.

1        16. The composition according to claim 1, wherein said  
2 peptide comprises at least one  $\alpha$ -amino acid.

1        17. The composition according to claim 1, wherein said  
2 peptide is formed from one or more amino acids selected from  
3 the group consisting of alanine, arginine, asparagine,  
4 aspartic acid, citrulline, cysteine, cystine, glutamine,  
5 glycine, histidine, isoleucine, leucine, lysine, methionine,  
6 ornithine, phenylalanine, phenylglycine, proline, serine,  
7 threonine, tryptophan tyrosine, valine, hydroxy proline,  
8  $\gamma$ -carboxyglutamate, O-phosphoserine,  $\beta$ -alanine,  $\alpha$ -amino  
9 butyric acid,  $\gamma$ -amino butyric acid,  $\alpha$ -amino isobutyric acid,  
10  $\epsilon$ -amino caproic acid, 7-amino heptanoic acid,  $\beta$ -aspartic  
11 acid,  $\gamma$ -glutamic acid, cysteine (ACM),  $\epsilon$ -lysine,  $\epsilon$ -lysine  
12 (A-Fmoc), methionine sulfone, norleucine, norvaline,  
13 ornithine, d-ornithine, p-nitro-phenylalanine, hydroxy  
14 proline, and thioproline.

1        18. The composition according to claim 17, wherein said  
2 peptide is formed from one or more amino acids selected from  
3 the group consisting of arginine, leucine, lysine,  
4 phenylalanine, tyrosine, valine, and phenylglycine.

1        19. The composition according to claim 1, wherein said  
2 acylated aldehyde or acylated ketone is acylated by an

3  
4 acylating agent having the formula  $R^9-\overset{\overset{O}{\parallel}}{C}-X$  wherein  $R^9$  is  
5 alkyl, alkenyl, cycloalkyl or aromatic, and X is a leaving  
6 group.

1       20. A dosage unit form comprising  
2           (A) a composition according to claim 1; and  
3           (B)   (a) an excipient,  
4                   (b) a diluent,  
5                   (c) a disintegrant,  
6                   (d) a lubricant,  
7                   (e) a plasticizer,  
8                   (f) a colorant,  
9                   (g) a dosing vehicle, or  
10                  (h) any combination thereof.

1       21. A dosage unit form according to claim 20 comprising  
2 a tablet, a capsule, or a liquid.

1       22. A method for administering a biologically-active  
2 agent to an animal in need of said agent, said method  
3 comprising administering orally to said animal a composition  
4 as defined in claim 1.

1       23. A method for preparing a composition, said method  
2 comprising mixing:  
3       (A) at least one biologically-active agent;  
4       (B) a carrier comprising  
5           (a)   (i) at least one acylated aldehyde of an  
6                   amino acid,  
7                   (ii) at least one acylated ketone of an amino  
8                   acid,  
9                   (iii) at least one acylated aldehyde of a  
10                  peptide,  
11                  (iv) at least one acylated ketone of a  
12                  peptide, or  
13                  (v) any combination of (a)(i), (a)(ii),  
14                  (a)(iii) and (a)(iv);

- 15 (b) (i) carboxymethyl-phenylalanine-leucine,  
 16 (ii) 2-carboxy-3-phenylpropionyl-leucine,  
 17 (iii) 2-benzylsuccinic acid, or  
 18 (iv) an actinonin, or  
 19 (v) a compound having the formula:



21 wherein: Ar is a substituted or  
 22 unsubstituted phenyl or naphthyl;

23 Y is  $\text{—}\overset{\text{O}}{\parallel}\text{C—}$  or  $\text{—SO}_2\text{—}$ ;  
 24

25 R<sup>1</sup> is  $\text{—N(R}^4\text{)—R}^3\text{—}\overset{\text{O}}{\parallel}\text{C—}$ ; wherein:  
 26

27 R<sup>3</sup> is C<sub>1</sub> to C<sub>24</sub> alkyl, C<sub>1</sub> to C<sub>24</sub> alkenyl,  
 28 phenyl, naphthyl, (C<sub>1</sub> to C<sub>10</sub> alkyl) phenyl, (C<sub>1</sub> to C<sub>10</sub>  
 29 alkenyl) phenyl, (C<sub>1</sub> to C<sub>10</sub> alkyl) naphthyl, (C<sub>1</sub> to C<sub>10</sub>  
 30 alkenyl) naphthyl, phenyl (C<sub>1</sub> to C<sub>10</sub> alkyl), phenyl (C<sub>1</sub>  
 31 to C<sub>10</sub> alkenyl), naphthyl (C<sub>1</sub> to C<sub>10</sub> alkyl) and naphthyl  
 32 (C<sub>1</sub> to C<sub>10</sub> alkenyl);

33 R<sup>3</sup> is optionally substituted with C<sub>1</sub> to C<sub>4</sub>  
 34 alkyl, C<sub>1</sub> to C<sub>4</sub> alkenyl, C<sub>1</sub> to C<sub>4</sub> alkoxy, -OH, -SH -CO<sub>2</sub>R<sup>5</sup>,  
 35 cycloalkyl, cycloalkenyl, heterocyclic, aryl, alkaryl,  
 36 heteroaryl or heteroalkaryl or any combination thereof;

37 R<sup>5</sup> is hydrogen, C<sub>1</sub> to C<sub>4</sub> alkyl or C<sub>1</sub> to C<sub>4</sub>  
 38 alkenyl;

39 R<sup>3</sup> is optionally interrupted by oxygen,  
 40 nitrogen, sulfur or any combination thereof; and

41 R<sup>4</sup> is hydrogen, C<sub>1</sub> to C<sub>4</sub> alkyl or C<sub>1</sub> to C<sub>4</sub>  
 42 alkenyl;

43 and n is from 1 to 5;

44 (vi) or any combination of (b) (i), (b) (ii),  
 45 (b) (iii), (b) (iv) and (b) (v);

46 (c) a combination of (a) and (b); and

47 (C) optionally a dosing vehicle.

48 24. A pharmacological composition comprising:

49 (A) at least one biologically-active agent; and

50 (B) at least one carrier comprising

- 51           (a)   (i) at least one acylated aldehyde of an  
52                   amino acid,  
53               (ii) at least one acylated ketone of an amino  
54                   acid,  
55               (iii) at least one acylated aldehyde of a  
56                   peptide,  
57               (iv) at least one acylated ketone of a  
58                   peptide, or  
59               (v) any combination of (a) (i), (a) (ii),  
60                   (a) (iii) and (a) (iv);
- 61           (b)   (i) carboxymethyl-phenylalanine-leucine,  
62                   (ii) 2-carboxy-3-phenylpropionyl-leucine,  
63                   (iii) 2-benzylsuccinic acid, or  
64                   (iv) 4-(phenylsulfonamido)-4-phenylbutyric  
65                   acid,  
66               (v) or any combination of (b) (i), (b) (ii),  
67                   (b) (iii) and (b) (iv); or  
68           (c)   a combination of (a) and (b).

1           25. The composition according to claim 24, wherein said  
2   biologically-active agent comprises at least one peptide,  
3   mucopolysaccharide, carbohydrate, or lipid.

1           26. The composition according to claim 25, wherein said  
2   biologically-active agent is selected from the group  
3   consisting of human growth hormone, bovine growth hormone,  
4   growth hormone-releasing hormone, an interferon,  
5   interleukin-II, insulin, heparin, calcitonin,  
6   erythropoietin, atrial natriuretic factor, an antigen, a  
7   monoclonal antibody, somatostatin, adrenocorticotropin,  
8   gonadotropin releasing hormone, oxytocin, vasopressin,  
9   vancomycin, desferrioxamine (DFO), or any combination  
10   thereof.

1        27. The composition according to claim 26, wherein said  
2 biologically-active agent comprises an interferon,  
3 interleukin-II, insulin, heparin, calcitonin, oxytocin,  
4 vasopressin, vancomycin, DFO and combinations thereof.

1        28. The composition according to claim 27, wherein said  
2 biologically-active agent comprises calcitonin.

1        29. The composition according to claim 24, wherein said  
2 amino acid is a naturally occurring amino acid.

1        30. The composition according to claim 24, wherein said  
2 amino acid is a synthetic amino acid.

1        31. The composition according to claim 24, wherein said  
2 amino acid is an  $\alpha$ -amino acid.

1        32. The composition according to claim 24, wherein said  
2 amino acid is selected from the group consisting of alanine,  
3 arginine, asparagine, aspartic acid, citrulline, cysteine,  
4 cystine, glutamine, glycine, histidine, isoleucine, leucine,  
5 lysine, methionine, ornithine, phenylalanine, phenylglycine,  
6 proline, serine, threonine, tryptophan tyrosine, valine,  
7 hydroxy proline,  $\gamma$ -carboxyglutamate, O-phosphoserine,  $\beta$ -  
8 alanine,  $\alpha$ -amino butyric acid,  $\gamma$ -amino butyric acid,  $\alpha$ -amino  
9 isobutyric acid,  $\epsilon$ -amino caproic acid, 7-amino heptanoic  
10 acid,  $\beta$ -aspartic acid,  $\gamma$ -glutamic acid, cysteine (ACM),  $\epsilon$ -  
11 lysine,  $\epsilon$ -lysine (A-Fmoc), methionine sulfone, norleucine,  
12 norvaline, ornithine, d-ornithine, p-nitro-phenylalanine,  
13 hydroxy proline, and thioproline.

1        33. The composition according to claim 32, wherein said  
2 amino acid is selected from the group consisting of  
3 arginine, leucine, lysine, phenylalanine, tyrosine, valine,  
4 and phenylglycine.

1        34. The composition according to claim 24, wherein said  
2 peptide is selected from the group consisting of a  
3 di-peptide, a tri-peptide, a tetra-peptide, or a  
4 penta-peptide.

1        35. The composition according to claim 24, wherein said  
2 peptide comprises at least one naturally occurring amino  
3 acid.

1        36. The composition according to claim 24, wherein said  
2 peptide comprises at least one synthetic amino acid.

1        37. The composition according to claim 24, wherein said  
2 peptide comprises at least one  $\alpha$ -amino acid.

1        38. The composition according to claim 24, wherein said  
2 peptide is formed from one or more amino acids selected from  
3 the group consisting of alanine, arginine, asparagine,  
4 aspartic acid, citrulline, cysteine, cystine, glutamine,  
5 glycine, histidine, isoleucine, leucine, lysine, methionine,  
6 ornithine, phenylalanine, phenylglycine, proline, serine,  
7 threonine, tryptophan tyrosine, valine, hydroxy proline,  
8  $\gamma$ -carboxyglutamate, O-phosphoserine,  $\beta$ -alanine,  $\alpha$ -amino  
9 butyric acid,  $\gamma$ -amino butyric acid,  $\alpha$ -amino isobutyric acid,  
10  $\epsilon$ -amino caproic acid, 7-amino heptanoic acid,  $\beta$ -aspartic  
11 acid,  $\gamma$ -glutamic acid, cysteine (ACM),  $\epsilon$ -lysine,  $\epsilon$ -lysine  
12 (A-Fmoc), methionine sulfone, norleucine, norvaline,  
13 ornithine, d-ornithine, p-nitro-phenylalanine, hydroxy  
14 proline, and thioproline.

1        39. The composition according to claim 38, wherein said  
2 peptide is formed from one or more amino acids selected from  
3 the group consisting of arginine, leucine, lysine,  
4 phenylalanine, tyrosine, valine, and phenylglycine.

1           40. The composition according to claim 24, wherein said  
2 acylated aldehyde or acylated ketone is acylated by an  
3  
4 acylating agent having the formula  $R^7-\overset{\overset{O}{\parallel}}{C}-X$  wherein  $R^7$  is  
5 alkyl, cycloalkyl, or aryl, and X is a leaving group.

1           41. The composition according to claim 40, wherein  $R^7$   
2 is methyl, ethyl, cyclohexane, cyclopentane, phenyl or  
3 benzyl.

1           42. The composition according to claim 40, wherein  $R^7$   
2 is cyclohexyl, cyclopentyl, or cycloheptyl, or acetyl.

1           43. A composition according to claim 24, wherein said  
2 biologically active agent comprises calcitonin, and said  
3 carrier comprises acetyl phenylalanine aldehyde and  
4 carbomethoxy phenylalanyl-leucine.

1           44. A composition according to claim 24, wherein said  
2 biologically-active agent comprises 1.5  $\mu\text{g/mL}$  of calcitonin,  
3 132 mg/ml of acetyl phenylalanine aldehyde, 33 mg/mL of  
4 carbomethoxy phenylalanyl-leucine and 25 mg/mL of acetyl-  
5 Phe-Leu-Leu-Arg aldehyde.

1           45. A dosage unit form comprising  
2           (A) a pharmacological composition according to  
3 claim 1; and  
4           (B) (a) an excipient,  
5               (b) a diluent,  
6               (c) a disintegrant,  
7               (d) a lubricant,  
8               (e) a plasticizer,  
9               (f) a colorant,  
10              (g) a dosing vehicle, or  
11              (h) any combination thereof.

1           46. A dosage unit form according to claim 45 comprising  
2 a tablet, a capsule, or a liquid.



1        47. A dosage unit form according to claim 45, wherein  
2 said dosing vehicle is selected from the group consisting of  
3 water, 1,2-propane diol, ethanol or any combination thereof.

1        48. A method for administering a biologically-active  
2 agent to a animal in need of said agent, said method  
3 comprising administering orally to said animal a composition  
4 as defined in claim 25.

1        49. A method for preparing a pharmacological  
2 composition, said method comprising mixing:

3        (A) at least one biologically-active agent;

4        (B) a carrier comprising

5            (a) (i) at least one acylated aldehyde of an  
6 amino acid,

7            (ii) at least one acylated ketone of an amino  
8 acid,

9            (iii) at least one acylated aldehyde of a  
10 peptide,

11           (iv) at least one acylated ketone of a  
12 peptide, or

13           (v) any combination of (a) (i), (a) (ii),  
14 (a) (iii) and (a) (iv);

15           (b) (i) carboxymethyl-phenylalanine-leucine,

16           (ii) 2-carboxy-3-phenylpropionyl-leucine,

17           (iii) 2-benzylsuccinic acid, or

18           (iv) 4-(phenylsulfonamido)-4-phenylbutyric  
19 acid,

20           (v) or any combination of (b) (i), (b) (ii),  
21 (b) (iii) and (b) (iv); or

22           (c) a combination of (a) and (b); and

23        (C) optionally a dosing vehicle.

1        50. The compound N-cyclohexanoylphenylalanine aldehyde.

1        51. A composition comprising an active agent and one or  
2 more modified amino acids comprising an amino acid amide or  
3 sulfonamide.

1        52. The composition according to claim 51, wherein said  
2 modified amino acids comprise a mixture of two or more amino  
3 acid amides or sulfonamides.

1        53. The composition according to claim 51, wherein said  
2 modified amino acids are derived from reaction of an amino  
3 acid with an amino reactive modifying agent comprises  
4 benzene sulfonyl chloride, benzoyl chloride, hippuryl  
5 chloride or amino acid carbodiimide.

1        54. The composition according to claim 51, wherein said  
2 biologically-active agent comprises human or bovine growth  
3 hormone, interferon, interleukin-II, insulin, calcitonin,  
4 erythropoietin, atrial natriuretic factor, antigens, or  
5 monoclonal antibody.

1        55. A composition comprising an active agent  
2 encapsulated within a modified amino acid microsphere, said  
3 microsphere comprising a wall of at least one amino acid  
4 amide or sulfonamide.

1        56. The composition according to claim 55, wherein said  
2 amino acid amide or sulfonamide comprise a benzene  
3 sulfonamide or benzene amide of an amino acid.

1        57. The composition according to claim 55, wherein said  
2 biologically-active agent comprises human or bovine growth  
3 hormone, interferon, interleukin-II, insulin, calcitonin,  
4 erythropoietin, atrial natriuretic factor, antigens, or  
5 monoclonal antibody.

1        58. A method for preparing a modified amino acid micro-  
2 sphere comprising the steps of:

- 3 (a) incubating at least one or more amino  
4 acid amide or sulfonamide in a aqueous acid solution so as  
5 to form said modified amino acid microsphere; and  
6 (b) collecting said modified amino acid  
7 microsphere.


1 59. The method according to claim 58, wherein said  
2 amino acid amide or sulfonamide are derived from reaction of  
3 an amino acid with an amino reactive modifying agent  
4 comprising benzene sulfonyl chloride, benzoyl chloride,  
5 hippuryl chloride or amino acid carbodiimide.

1 60. The method according to claim 58, wherein said  
2 aqueous acid solution further comprises a biologically-  
3 active agent.


1 61. The method according to claim 60, wherein said  
2 biologically-active agent comprises human or bovine growth  
3 hormone, interferon, interleukin-II, insulin, calcitonin,  
4 erythropoietin, atrial naturetic factor, antigens, or  
5 monoclonal antibody.

1 62. The composition according to claim 1, wherein said  
2 amino acid or peptide is acylated or sulfonated by an  
3  
4 acylating agent having the formula  $R^7-\overset{\overset{O}{\parallel}}{C}-X$  or a sulfonating  
5 agent having the formula  $R^7-SO_2-X$  wherein  $R^7$  is alkyl,  
6 alkenyl, or aromatic, and X is a leaving group.


1 / 22



2 / 22



3 / 22








FIG. 4

5 / 22





6 / 22







FIG. 7

8 / 22


FIG. 8




SUBSTITUTE SHEET (RULE 26)


9 / 22

FIG. 9






11 / 22



12 / 22

FIG. 12



—▲— 3-(Phenylsulfonamido) benzoic acid 400mg/kg + sCT 10ug/kg  
—▼— 4-(Phenylsulfonamido) hippuric acid 400mg/kg + sCT 10ug/kg

13 / 22

FIG. 13










FIG. 14










17 / 22







20 / 22







FIG. 21





## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/04560

**A. CLASSIFICATION OF SUBJECT MATTER**

IPC(5) : Please See Extra Sheet.

US CL : 424/490

According to International Patent Classification (IPC) or to both national classification and IPC

**B. FIELDS SEARCHED**

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/490, 491, 489

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

**C. DOCUMENTS CONSIDERED TO BE RELEVANT**

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, A, 4,835,312 (ITOH ET AL) 30 May 1989, see entire document.	1-62
Y	US, A, 4,976,968 (STEINER) 11 December 1990, see entire document.	1-62



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T	later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*A* document defining the general state of the art which is not considered to be part of particular relevance	*X*	document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*E* earlier document published on or after the international filing date	*Y*	document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Z*	document member of the same patent family
*O* document referring to an oral disclosure, use, exhibition or other means		
*P* document published prior to the international filing date but later than the priority date claimed		

Date of the actual completion of the international search

17 JULY 1994

Date of mailing of the international search report

16 AUG 1994

Name and mailing address of the ISA/US  
Commissioner of Patents and Trademarks  
Box PCT  
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

WILLIAM BENSTON, JR.

Telephone No. (703) 308-2351

**INTERNATIONAL SEARCH REPORT**

International application No.  
PCT/US94/04560

**A. CLASSIFICATION OF SUBJECT MATTER:**  
**IPC (5):**

A61L 9/16

**This Page is Inserted by IFW Indexing and Scanning  
Operations and is not part of the Official Record**

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☐ BLACK BORDERS
- ☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES
- ☐ FADED TEXT OR DRAWING
- ☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING
- ☐ SKEWED/SLANTED IMAGES
- ☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS
- ☐ GRAY SCALE DOCUMENTS
- ☒ LINES OR MARKS ON ORIGINAL DOCUMENT
- ☐ REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
- ☐ OTHER: \_\_\_\_\_

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.**